



Checklist for Actemra (tocilizumab) Referral

Patient Name: _____ DOB: _____ Date: _____

Referring Physician: _____ NPI: _____

Office Contact/Title/Email: _____

Office Address: _____

Office Phone: _____ Office Fax: _____

Best contact number for physician in case of reaction: _____

Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
 - Note should include any therapies tried/failed, and must include discussion about Actemra
 - Medication list and allergies
- Supporting lab reports/imaging for Actemra treatment
 - Baseline: Lipid panel, LFTs, neutrophils, platelets, and negative TB screening
 - *Recommended:* negative hepatitis B screening
 - For continued therapy:
 - Lipid panel 4-8 weeks after Actemra start
 - LFTs every 4-8 weeks for first 6 months after Actemra start, then every 3 months thereafter
 - Neutrophils and platelets 4-8 weeks after Actemra start, then every 3 months thereafter
- Actemra Prescribing Order (see attached)

-- We will obtain prior authorization and schedule your patient as soon as possible --

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 ext. 112

Visit our website: www.InfusionCenterNE.com
Email: info@infusioncenterne.com

Infusion Center of New England
9 Payson Road, Suite 100, Foxboro MA 02035
Ph: 781-551-5812
Fax: 508-698-8671



Prescribing Order: Actemra (tocilizumab)

Date of Order: _____

New Start Maintenance

Date of last infusion: _____

Patient Name: _____ DOB: _____ M/F: _____

Diagnosis (include ICD-10 code/s): _____

NKDA Allergies: _____

Patient Weight: _____

Premedication:

- | | |
|--|--|
| <input type="checkbox"/> Acetaminophen 1000mg PO | <input type="checkbox"/> Diphenhydramine 25mg IV |
| <input type="checkbox"/> Loratadine 10mg or Cetirizine 10mg PO | <input type="checkbox"/> Solu-medrol 125mg IV |
| <input type="checkbox"/> Diphenhydramine 25mg PO | <input type="checkbox"/> Other: _____ |

Lab Orders:

- | | |
|--|---|
| <input type="checkbox"/> CBC w/diff, CMP every _____ weeks | <input type="checkbox"/> LFTs every _____ weeks |
| <input type="checkbox"/> Lipid Panel 4-8 weeks after treatment start x 1 | |
| <input type="checkbox"/> Other: _____ | |

Actemra Medication Order

Dosing: 4mg/kg 8mg/kg Other: _____

Frequency: Every 2 weeks Every 4 weeks Other: _____

Administration:

- ✓ Mix Actemra in 100ml Normal Saline and administer intravenously over 1 hour.
- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ HOLD Actemra if most recent ANC < 2000, AST/ALT > 1.5x normal limit, or platelets <100,000. Notify physician.
- ✓ In case of infusion reaction, STOP infusion and follow NCNE infusion reaction protocol. Notify physician.

Ordering Provider Name

NPI

Signature

Date