

## **Checklist for Infliximab Referral**

Patient	t Name:		DOB:		Date:		
Referri	Referring Physician: NPI: NPI:						
Referri	ing Office Contact/Title	e:					
Office A	Address:						
Office Phone: Office Fax:							
Best co	ontact number for phy	sician in case of reactio	n:				
Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.							
	□ Patient demographic information						
	☐ Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance						
	Supporting clinical notes and office visits. Two notes preferred.						
		nclude any therapies tr st and allergies	ied/failed, and m	ust include discus	ssion about Infliximat	0	
	Supporting lab reports/orders for Infliximab treatment						
_	<ul> <li>Baseline: CBC w diff, LFTs, negative Hepatitis B screening, and negative TB screening</li> <li>Recommended: monitor CBC w diff and LFTs periodically with treatment</li> <li>Infliximab Prescribing Order (see attached)</li> </ul>						
	We will ob	otain prior authorizat	ion and schedu	le your patient	as soon as possible	: <del></del>	

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 ext. 112

Visit our website: <a href="www.InfusionCenterNE.com">www.InfusionCenterNE.com</a>
Email: info@infusioncenterne.com



## **Prescribing Order: Infliximab**

Date of Order:	☐ New Start ☐ Maintenance  Date of last infusion:							
Patient Name:	DOB: M/F:							
Diagnosis (include ICD-10 code/s):								
□ NKDA Allergies:								
Patient Weight:								
Premedication:								
☐ Acetaminophen 1000mg PO	Diphenhydramine 25mg IV							
☐ Loratadine 10mg or Cetirizine 10mg PO	☐ Solu-medrol 125mg IV							
☐ Diphenhydramine 25mg PO	☐ Other:							
Lab Orders:  ☐ CBC w/diff, CMP every weeks	☐ LFTs every weeks							
□ Other:								
Infliximab Medication Order □ Influse Remicade OR □ Infliximab biosimilar as required by patient's insurance Dosing: □ 5mg/kg □ 10mg/kg □ Other:								
Frequency:	very 8 weeks							
Administration:  ✓ Reconstitute each 100 mg Infliximab vial with 10 mL of Sterile Water for Injection.  ✓ Determine dose based on patient weight then dilute the required volume of the reconstituted Infliximab solution in 250 mL 0.9% Sodium Chloride.  ✓ Infuse Infliximab over at least 2 hours with in-line 0.2 micron filter.  ✓ Do not administer if patient has active signs or symptoms of infection.  ✓ In case of infusion reaction, STOP infusion and follow NCNE infusion reaction protocol. Notify physician.								
Ordering Provider Name	NPI							
Signature	Date							