



**Checklist for Stelara (ustekinumab) Referral**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Date: \_\_\_\_\_

Referring Physician: \_\_\_\_\_ NPI: \_\_\_\_\_

Office Contact/Title/Email: \_\_\_\_\_

Office Address: \_\_\_\_\_

Office Phone: \_\_\_\_\_ Office Fax: \_\_\_\_\_

Best contact number for physician in case of reaction: \_\_\_\_\_

*Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.*

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). \*Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
  - Note should include any therapies tried/failed, and must include discussion about Stelara
  - Medication list and allergies
- Supporting lab reports/orders for Stelara treatment
  - Baseline: negative TB screening
  - *Recommended:* baseline CBC w diff and CMP
- Stelara Prescribing Order (see attached)

**-- We will obtain prior authorization and schedule your patient as soon as possible --**

---

**Fax all information to our Infusion Coordinator: 508-698-8671**

Call with any questions: 781-551-5812 ext. 112

---

Visit our website: [www.InfusionCenterNE.com](http://www.InfusionCenterNE.com)

Email: [info@infusioncenterne.com](mailto:info@infusioncenterne.com)

**Infusion Center of New England**

9 Payson Road, Suite 100, Foxboro MA 02035

Ph: 781-551-5812

Fax: 508-698-8671



**Prescribing Order: Stelara (ustekinumab)**

Date of Order: \_\_\_\_\_

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ M/F: \_\_\_\_\_

Diagnosis (include ICD-10 code/s): \_\_\_\_\_

NKDA Allergies: \_\_\_\_\_

Patient Weight: \_\_\_\_\_

**Premedication:**

- |                                                                |                                                  |
|----------------------------------------------------------------|--------------------------------------------------|
| <input type="checkbox"/> Acetaminophen 1000mg PO               | <input type="checkbox"/> Diphenhydramine 25mg IV |
| <input type="checkbox"/> Loratadine 10mg or Cetirizine 10mg PO | <input type="checkbox"/> Solu-medrol 125mg IV    |
| <input type="checkbox"/> Diphenhydramine 25mg PO               | <input type="checkbox"/> Other: _____            |

**Lab Orders:**

\_\_\_\_\_

**Stelara Medication Order**

- Initial Induction Infusion: *Indicated for Crohn's Disease and Ulcerative Colitis*  
Dilute in 250ml 0.9% Sodium Chloride and administer intravenously over 1 hour using 0.2 micron filter.
- 260mg (up to 55kg)
  - 390mg (55 - 85kg)
  - 520mg (>85kg)

**Administration:**

- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ In case of infusion reaction, STOP infusion and follow NCNE infusion reaction protocol. Notify physician.

---

Ordering Provider Name \_\_\_\_\_ NPI \_\_\_\_\_

---

Signature \_\_\_\_\_ Date \_\_\_\_\_