



Checklist for Rituximab Referral

Patient Name: _____ DOB: _____ Date: _____

Referring Physician: _____ NPI: _____

Referring Office Contact/Title: _____

Office Address: _____

Office Phone: _____ Office Fax: _____

Best contact number for physician in case of reaction: _____

Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
 - Note should include any therapies tried/failed, and must include discussion about Rituximab
 - Medication list and allergies
- Supporting lab reports/orders for Rituximab treatment
 - Baseline: CBC w diff, Plt, CMP, Hepatitis B surface antibody, Hepatitis B core antibody, and negative TB screening
 - Continued Therapy: monitor CBC w diff and platelets every 2-4 months
- Rituximab Prescribing Order (see attached)

-- We will obtain prior authorization and schedule your patient as soon as possible --

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4

Visit our website: www.InfusionCenterNE.com
Email: info@infusioncenterne.com

Infusion Center of New England
9 Payson Road, Suite 100, Foxboro MA 02035
Ph: 781-551-5812
Fax: 508-698-8671



Prescribing Order: Rituximab

Date of Order: _____

New Start

Maintenance

Date of last infusion: _____

Patient Name: _____ DOB: _____ M/F: _____

Diagnosis (include ICD-10 code/s): _____

NKDA Allergies: _____

Patient Weight: _____

Premedication:

Acetaminophen 1000mg PO

Loratadine 10mg or Cetirizine 10mg PO

Diphenhydramine 25mg PO

Diphenhydramine 25mg IV

Solu-medrol 125mg IV

Other: _____

Lab Orders:

CBC w/diff, Plt every _____ months

Other: _____

Infliximab Medication Order

Infuse Rituxan OR Rituximab biosimilar as required by patient's insurance

Dosing: 500mg 1000mg Other: _____

Frequency: On Day 1 and Day 15 every 6 months Other: _____

Administration:

- ✓ Dilute Rituximab in 0.9% Sodium Chloride, volume (ml) equal to Rituxan dose (mg)
- ✓ First infusion: Initiate infusion at rate of 50ml/hr and increase 50ml/hr every 30min to max rate of 400ml/hr
Subsequent Infusions: Initiate infusion at rate of 100ml/hr and increase 100ml/hr every 30min to a maximum rate of 400ml/hr
- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ In case of infusion reaction, STOP infusion and follow ICNE/NCNE infusion reaction protocol. Notify physician.

Ordering Provider Name

NPI

Signature

Date

Infusion Center of New England
9 Payson Road, Suite 100, Foxboro MA 02035
Ph: 781-551-5812
Fax: 508-698-8671