

Checklist for Vyvgart (efgartigimod alfa-fcab) Referral

Patient	nt Name: DOB: Date:					
Referri	Referring Physician: NPI: NPI:					
Office Contact/Title/Email:						
Office A	Address:					
Office I	Phone: Office Fax:					
Best co	ontact number for physician in case of reaction:					
Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.						
	☐ Patient demographic information					
	☐ Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance					
	☐ Supporting clinical notes and office visits. Two notes preferred.					
	 Note should include any therapies tried/failed, and must include discussion about Vyvgart Medication list and allergies 					
	☐ Supporting lab reports/testing for Vyvgart treatment					
	 Required: Positive serologic test for anti-AChR antibodies 					
	 EMG, nerve stimulation studies, positive anticholinesterase test 					
	Vyvgart Prescribing Order (see attached)					
	We will obtain prior authorization and schedule your patient as soon as possible					

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4

Visit our website: www.InfusionCenterNE.com
Email: info@infusioncenterNE.com



Prescribing Order: Vyvgart IV and Vyvgart Hytrulo

Date of Order:						
Patient Nar	me:	DOB:	M/F:			
Diagnosis (include ICD-10 code/s):						
□ NKDA	Allergies:					
Patient Wei	ight:					
Premedicat	ion:					
☐ Acetam	inophen 1000mg PO	Diphenhydramine 25	mg IV			
☐ Loratad	line 10mg or Cetirizine 10mg PO	☐ Solu-medrol 125mg IV				
☐ Diphen	hydramine 25mg PO	☐ Other:				
Lab Orders:	:					
Vyvgart IV	Medication Order					
Pat	ients < 120kg: Vyvgart 10mg/kg IV weekly	x4 weeks PRN				
Pat	ients ≥ 120kg: Vyvgart 1200mg IV weekly >	4 weeks PRN				
Ma						
√ Dilu	ite Vyvgart in 0.9% Sodium Chloride for a	r via 0.22 micron filter over 1				
hour						
Vyvgart Hytrulo Medication Order ☐ Vyvgart Hytrulo 1008mg SQ weekly x4 weeks PRN						
					Maintenance schedule based on clinical evaluation:	
✓ Inje	ect Vyvgart Hytrulo 1008mg subcutaneous	y into abdomen over 30-90 seconds				
Administra	tion:					
✓ Do	 ✓ Do not administer if patient has active signs or symptoms of infection. ✓ Monitor patient for 1 hour following IV administration, 30 minutes following SQ administration. 					
✓ Mo						
✓ In c	ase of infusion/injection reaction, STOP ar	nd follow NCNE infusion reaction pro	tocol. Notify physician.			
Ordering Pr	rovider Name	NPI				
Signature		Date				