

Checklist for Skyrizi (risankizumab-rzaa) Referral

Patient	ent Name: DO	B:	Date:				
Referri	erring Physician:	NPI:					
Office	ce Contact/Title/Email:						
Office A	ce Address:						
Office	ce Phone: Office	ce Fax:					
Best co	t contact number for physician in case of reaction:						
Please	se return completed checklist and checklist items to initio	ate referral. Use this	s form as fax cover sheet.				
	☐ Patient demographic information						
	☐ Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance						
	☐ Supporting clinical notes and office visits. Two notes preferred.						
	Note should include any therapies tried/faileMedication list and allergies	ed, and must include	e discussion about Skyrizi				
	☐ Supporting lab reports/imaging for Skyrizi IV Induction treatment						
	 Baseline negative TB screening 						
	 Baseline liver enzymes and bilirubin levels 						
	☐ Skyrizi Prescribing Order (see attached)						
	We will obtain prior authorization and	l schedule your pa	atient as soon as possible –				

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4

Visit our website: www.InfusionCenterNE.com
Email: info@infusioncenterne.com



Prescribing Order: Skyrizi (risankizumab-rzaa) IV Induction

Date of Ord	ler:		☐ New Start	☐ Maintenance Date of last injection:					
Patient Nar	ne:		DOB: _	M/F	M/F:				
Diagnosis (include ICD-10 code/s):									
□ NKDA	All	lergies:							
Premedicat	_								
Acetam	inophe	n 1000mg PO	Diphe	nhydramine 25mg PO					
☐ Loratad	ine 10n	ng or Cetirizine 10mg PO	☐ Other	·					
Skyrizi Med	lication	Order – Induction Dosing							
		Skyrizi 600mg IV over 1 hour (Crohn's disease)						
Dosing:		☐ Skyrizi 1200mg IV over 2 hours (ulcerative colitis)							
Frequency:		Week 0, Week 4, and Week 8							
Administra	tion:								
✓ Adr	ninister	Skyrizi IV in 250ml 5% Dextros	e or 0.9% Sodium Chlorid	e over 1-2 hours per indication					
✓ Do not administer if patient has active signs or symptoms of infection or hepatic dysfunction.									
✓ In case of reaction, follow ICNE infusion reaction protocol. Notify physician.									
Ordering Provider Name			NI	PI					
Signature			D	ate	_				
Signature			Da	are					