

Checklist for Kisunla (donanemab) Referral

Patien	nt Name:	: Dob: Dob: Date:			
Referring Physician:		sician:NPI:NPI:			
Office	Contact,	:/Title/Email:			
Office	Address	s:			
Office	Phone:	Office Fax:			
Best c	ontact n	number for physician in case of reaction:			
Please	return c	completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.			
	Patien	t demographic information			
	Insura	nce information and copy of insurance card/s (front and back). *Include primary and secondary insuranc			
	Suppo	Supporting clinical notes and office visits. Two notes preferred.			
	0	Note should include any therapies tried/failed, and must include discussion about Kisunla Medication list and allergies			
	o Suppo	Cognitive assessment and functional assessment with score and interpretation orting lab reports/imaging for Kisunla treatment			
	0	MRI within 1 year of treatment start			
	0	Confirmation of amyloid beta pathology (LP or PET Scan)			
	0	ApoE testing to determine ARIA risk			
	0	CMS Registration (must be completed every 6 months)			
	Durabl	le Power of Attorney for Health Care (DPAHC), if applicable			
	Kisunla	a Prescribing Order and Indication Checklist (see attached)			

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4



Prescribing Order: Kisunla (donanemab)

Date of Order:	☐ New Start ☐ Maintenance Date of last infusion:					
Patient Name:	DOB:	M/F:				
Diagnosis (include ICD-10 code/s):						
□ NKDA Allergies:						
Patient Weight:	_					
Premedication: ☐ Acetaminophen 1000mg PO ☐ Loratadine 10mg or Cetirizine 10mg PO ☐ Diphenhydramine 25mg PO Lab Orders:	☐ Diphenhydramine 25 ☐ Solu-medrol 125mg IV ☐ Other:	•				
_						
■ Infusion 1: 350m ■ Infusion 2: 700m ■ Infusion 3: 1050r ■ Infusion 4 and beyon □ Maintenance: Kisunla 1400mg every	g mg d: 1400mg					
 Administration: ✓ Hold infusion if no MRI Brain prior to the 2nd, 3rd, 4th and 7th infusion ✓ Hold infusion and notify provider if patient experiencing any of the following signs of ARIA: ○ Headache, Confusion, Dizziness, Nausea, Vision Changes, Gait Changes, Seizures ✓ In case of infusion reaction, STOP infusion and follow ICNE infusion reaction protocol. Notify physician. 						
Ordering Provider Name	NPI					
Signature	Date					



Patient Name:	DOE	3:
i aticiit ivaiiic.		'·

Legembi/Kisunla Indication Checklist

INFORMATION REQUIRED FOR TREATMENT INITIATION				
1)	Patient ICD-10 (select all that apply)	Clinical Diagnosis (select one)		
	G30.0 Alzheimer's disease, early onset	☐ Mild cognitive impairment due to AD		
	☐ G30.1 Alzheimer's disease, late onset	☐ Mild AD Dementia		
	☐ G30.9 Alzheimer's disease, unspecified			
	☐ G31.84 Mild cognitive impairment			
2)	Cognitive Screening: within 6 months Date:			
	Name of Test Used:	Score:		
	Interpretation:			
	Functional Screening: within 6 months Date:			
	Name of Test Used:	Score:		
3)	Confirmation of Amyloid-Beta Pathology: MUST provide supporting documentation			
	☐ Amyloid PET Scan Date:	Result:		
	☐ CSF Amyloid Confirmation Date:	Result:		
4)	Monitoring for Amyloid Polated Imaging Ahno	ormalities (ARIA)		
4)	Monitoring for Amyloid Related Imaging Abnormalities (ARIA) ***LEQEMBI/KISUNLA requires brain MRI within 1 year of treatment start date***			
	☐ Initial MRI Brain Date:			
	Evidence of ARIA-E Negative	□ Positive		
	Evidence of ARIA-H 🔲 Negative			
5)	Schedule for MRI Monitoring: the following is	required by Infusion Center of New England		
3)	☐ Legembi: Obtain MRI prior to the 5 th , 7			
	☐ Kisunla: Obtain MRI prior to the 2 nd , 3 rd	•		
6)	Is the patient on anticoagulation? Yes			
•	Is the patient on antiplatelets?			
	Has ApoE testing been performed? Yes			
0)	That Apol testing occur performed: 103	Result:		
9)	CMS REGISTRATION NUMBER			
	(REQUIRED)	Date:		