



Checklist for EVENITY or PROLIA Referral

Patient Name: _____ DOB: _____ Date: _____

Referring Physician: _____ NPI: _____

Office Contact/Title/Email: _____

Office Address: _____

Office Phone: _____ Office Fax: _____

Best contact number for physician in case of reaction: _____

Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
 - Note should include any therapies tried/failed, and must include discussion about Evenity / Prolia
 - Medication list and allergies
 - All patients should be supplemented with calcium 1000 mg daily and at least 400 IU vitamin D daily**
- Supporting lab reports/imaging for Evenity / Prolia treatment
 - Serum calcium and vitamin D levels at baseline and continual monitoring
 - Serum evaluation of kidney function at baseline and continual monitoring: BUN/Cr, GFR
- Evenity / Prolia Prescribing Order (see attached)

-- We will obtain prior authorization and schedule your patient as soon as possible --

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4

Visit our website: www.InfusionCenterNE.com

Email: info@infusioncenterne.com

Infusion Center of New England
18 Washington Street, Foxboro MA 02035
Ph: 781-551-5812
Fax: 508-698-8671



Prescribing Order: Evenity (romosozumab) / Prolia (denosumab)

Date of Order: _____

New Start Maintenance

Date of last injection: _____

Patient Name: _____

DOB: _____

M/F: _____

Diagnosis (include ICD-10 code/s): _____

NKDA Allergies: _____

Premedication:

Acetaminophen 1000mg PO
 Loratadine 10mg or Cetirizine 10mg PO

Diphenhydramine 25mg PO
 Other: _____

Medication Order

EVENITY 105mg/1.17ml x2 - Total dose 210mg

Monthly x 12 months

Other: _____

PROLIA 60mg/1ml (or biosimilar as required by patient's insurance)

Every 6 months

Other: _____

Administration:

- ✓ Administer Evenity / Prolia (or biosimilar) as a subcutaneous injection/s in the thigh, abdomen, or upper arm
- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ In case of reaction, follow NCNE infusion reaction protocol. Notify physician.

Ordering Provider Name

NPI

Signature

Date