



## Checklist for Simponi Aria (golimumab) Referral

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Date: \_\_\_\_\_

Referring Physician: \_\_\_\_\_ NPI: \_\_\_\_\_

Referring Office Contact/Title: \_\_\_\_\_

Office Address: \_\_\_\_\_

Office Phone: \_\_\_\_\_ Office Fax: \_\_\_\_\_

Best contact number for physician in case of reaction: \_\_\_\_\_

*Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.*

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). \*Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
  - Note should include any therapies tried/failed, and must include discussion about Simponi Aria
  - Medication list and allergies
- Supporting lab reports/orders for Simponi Aria treatment
  - Required Baseline:* TB screening, HBV screening, CBC w diff
  - Recommended:* periodic CBC w diff
- Simponi Aria Prescribing Order (see attached)

**-- We will obtain prior authorization and schedule your patient as soon as possible --**

---

**Fax all information to our Infusion Coordinator: 508-698-8671**

Call with any questions: 781-551-5812 option 4

---

Visit our website: [www.InfusionCenterNE.com](http://www.InfusionCenterNE.com)

Email: [info@infusioncenterne.com](mailto:info@infusioncenterne.com)

Infusion Center of New England  
18 Washington Street, Foxboro MA 02035  
Ph: 781-551-5812  
Fax: 508-698-8671



### Prescribing Order: Simponi Aria (golimumab)

Date of Order: \_\_\_\_\_

New Start  Maintenance

Date of last infusion: \_\_\_\_\_

Patient Name: \_\_\_\_\_

DOB: \_\_\_\_\_

M/F: \_\_\_\_\_

Diagnosis (include ICD-10 code/s): \_\_\_\_\_

NKDA Allergies: \_\_\_\_\_

Patient Weight: \_\_\_\_\_

#### Premedication:

Acetaminophen 1000mg PO  
 Loratadine 10mg or Cetirizine 10mg PO  
 Diphenhydramine 25mg PO

Diphenhydramine 25mg IV  
 Solu-medrol 125mg IV  
 Other: \_\_\_\_\_

#### Lab Orders:

CBC w/diff every \_\_\_\_\_ weeks  CMP every \_\_\_\_\_ weeks  
 Other: \_\_\_\_\_

#### Medication Order

Dosing:  Simponi Aria 2mg/kg IV

Frequency:  Dose at week 0, 4 and 8  Maintenance dose every 8 weeks thereafter

#### Administration:

- ✓ Dilute Simponi Aria in 100ml 0.9% Sodium Chloride. Administer intravenously over 30 min
- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ Do not administer in patient with signs/symptoms of CHF, TB, or hepatitis
- ✓ In case of infusion reaction, STOP infusion and follow ICNE infusion reaction protocol. Notify physician.

---

Ordering Provider Name

NPI

---

Signature

Date