



Checklist for Stelara (ustekinumab) Referral

Patient Name: _____ DOB: _____ Date: _____

Referring Physician: _____ NPI: _____

Office Contact/Title/Email: _____

Office Address: _____

Office Phone: _____ Office Fax: _____

Best contact number for physician in case of reaction: _____

Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
 - Note should include any therapies tried/failed, and must include discussion about Stelara
 - Medication list and allergies
- Supporting lab reports/orders for Stelara treatment
 - Baseline: negative TB screening
 - Recommended: baseline CBC w diff and CMP
- Stelara Prescribing Order (see attached)

-- We will obtain prior authorization and schedule your patient as soon as possible --

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4

Visit our website: www.InfusionCenterNE.com

Email: info@infusioncenterne.com



Prescribing Order: Stelara (ustekinumab)

Date of Order: _____

Patient Name: _____ DOB: _____ M/F: _____

Diagnosis (include ICD-10 code/s): _____

NKDA Allergies: _____

Patient Weight: _____

Premedication:

<input type="checkbox"/> Acetaminophen 1000mg PO	<input type="checkbox"/> Diphenhydramine 25mg IV
<input type="checkbox"/> Loratadine 10mg or Cetirizine 10mg PO	<input type="checkbox"/> Solu-medrol 125mg IV
<input type="checkbox"/> Diphenhydramine 25mg PO	<input type="checkbox"/> Other: _____

Lab Orders:

Stelara Medication Order

Initial Induction Infusion: *Indicated for Crohn's Disease and Ulcerative Colitis*

Dilute in 250ml 0.9% Sodium Chloride and administer intravenously over 1 hour using 0.2 micron filter.

- 260mg (up to 55kg)
- 390mg (55 - 85kg)
- 520mg (>85kg)

Administration:

- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ In case of infusion reaction, STOP infusion and follow NCNE infusion reaction protocol. Notify physician.

Ordering Provider Name

NPI

Signature

Date