



### **Checklist for Tysabri (natalizumab) Referral**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Date: \_\_\_\_\_

Referring Physician: \_\_\_\_\_ NPI: \_\_\_\_\_

Office Contact/Title/Email: \_\_\_\_\_

Office Address: \_\_\_\_\_

Office Phone: \_\_\_\_\_ Office Fax: \_\_\_\_\_

Best contact number for physician in case of reaction: \_\_\_\_\_

*Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.*

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). \*Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
  - Note should include any therapies tried/failed, and must include discussion about Tysabri
  - Medication list and allergies
- Supporting lab reports/imaging for Tysabri treatment
  - Baseline: MRI Brain, JCV antibody, CBC w diff, CMP, Varicella antibody
  - Recommended for continued therapy:* frequent MRI Brain, JCV, CBC w diff, and CMP monitoring
- Prescriber must be registered in the TOUCH® Prescribing Program to prescribe Tysabri
  - Provider must authorize continued treatment every 6 months
- Tysabri Prescribing Order (see attached)

**-- We will obtain prior authorization and schedule your patient as soon as possible --**

---

**Fax all information to our Infusion Coordinator: 508-698-8671**

Call with any questions: 781-551-5812 option 4

---

Visit our website: [www.InfusionCenterNE.com](http://www.InfusionCenterNE.com)

Email: [info@infusioncenterNE.com](mailto:info@infusioncenterNE.com)

Infusion Center of New England  
18 Washington Street, Foxboro MA 02035  
Ph: 781-551-5812  
Fax: 508-698-8671



### Prescribing Order: Tysabri (natalizumab)

Date of Order: \_\_\_\_\_

New Start  Maintenance

Date of last infusion: \_\_\_\_\_

Patient Name: \_\_\_\_\_

DOB: \_\_\_\_\_ M/F: \_\_\_\_\_

Diagnosis (include ICD-10 code/s): \_\_\_\_\_

NKDA Allergies: \_\_\_\_\_

Patient Weight: \_\_\_\_\_

#### Premedication:

- Acetaminophen 1000mg PO
- Loratadine 10mg or Cetirizine 10mg PO
- Diphenhydramine 25mg PO

- Diphenhydramine 25mg IV
- Solu-medrol 125mg IV
- Other: \_\_\_\_\_

#### Lab Orders:

- JCV antibody every \_\_\_\_\_ months
- CBC w diff, CMP every \_\_\_\_\_ months

#### Tysabri Medication Order

- Tysabri 300mg in 100ml NS IV over 1 hour every 4 weeks x 12 months
- Tysabri 300mg in 100ml NS IV over 1 hour every \_\_\_\_ weeks x 12 months
- Monitor patient for 1 hour after Tysabri infusion for first 12 treatments, then monitor per ICNE protocol. Infuse normal saline 100ml/hr during monitoring period.

#### Administration:

- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ Do not administer if any suspected signs/symptoms of PML.
- ✓ Complete required TOUCH screening and checklist prior to each infusion.
- ✓ In case of infusion reaction, STOP infusion and follow ICNE infusion reaction protocol. Notify physician.

---

Ordering Provider Name

NPI

---

Signature

Date