



Checklist for Ultomiris (ravulizumab) Referral

Patient Name: _____ DOB: _____ Date: _____

Referring Physician: _____ NPI: _____

Office Contact/Title/Email: _____

Office Address: _____

Office Phone: _____ Office Fax: _____

Best contact number for physician in case of reaction: _____

Please return completed checklist and checklist items to initiate referral. Use this form as fax cover sheet.

- Patient demographic information
- Insurance information and copy of insurance card/s (front and back). *Include primary and secondary insurance
- Supporting clinical notes and office visits. Two notes preferred.
 - Note should include any therapies tried/failed, and must include discussion about Ultomiris
 - Medication list and allergies
- Supporting lab reports/imaging for Ultomiris treatment
- Ultomiris Prescribing Order (see attached)

-- We will obtain prior authorization and schedule your patient as soon as possible --

Fax all information to our Infusion Coordinator: 508-698-8671

Call with any questions: 781-551-5812 option 4

Visit our website: www.InfusionCenterNE.com
Email: info@infusioncenterNE.com



Prescribing Order: Ultomiris (ravulizumab)

Date of Order: _____

Patient Name: _____ DOB: _____ M/F: _____

Diagnosis (include ICD-10 code/s): _____

NKDA Allergies: _____

Patient Weight: _____

Premedication:

<input type="checkbox"/> Acetaminophen 1000mg PO	<input type="checkbox"/> Diphenhydramine 25mg IV
<input type="checkbox"/> Loratadine 10mg or Cetirizine 10mg PO	<input type="checkbox"/> Solu-medrol 125mg IV
<input type="checkbox"/> Diphenhydramine 25mg PO	<input type="checkbox"/> Other: _____

Lab Orders:

Ultomiris Medication Order

✓ Ultomiris IV dosing per table:

Body Weight	Loading Dose	Maintenance Dose
40 – 60kg	2400mg	3000mg
60 – 100kg	2700mg	3300mg
>100kg	3000mg	3600mg

✓ Dilute Ultomiris in 0.9% Sodium Chloride for a final concentration of 5mg/ml and infuse via 0.22 micron filter at appropriate rate per weight per manufacturer medication guidelines

Frequency: Loading Dose Maintenance at week 2 and every 8 weeks
 Every 8 weeks

Administration:

- ✓ Do not administer if patient has active signs or symptoms of infection.
- ✓ In case of infusion reaction, STOP infusion and follow NCNE infusion reaction protocol. Notify physician.

Ordering Provider Name

NPI

Signature

Date